

REVIEW ARTICLE

Meditation and De-Addiction: A Neurobiological Perspective

KM Megha¹, Arjun Ram Roj¹, Vijay Shanker Yadav²,
Chaitanya Kaushal³, Pallabi Pal¹, Monalisa Das⁴, Sanjib Patra^{1*}

Megha KM, Roj AR, Yadav VS, et al. **Meditation and De-Addiction: A Neurobiological Perspective.** *Int J Endovasc Treat Innov Tech.* 2025;6(2):76-85.

Abstract

Background: Addiction is a multifaceted disorder involving compulsive substance use, impaired self-regulation, and continued use despite adverse consequences. Neurobiological alterations in key brain regions, including the Prefrontal Cortex (PFC), amygdala, hippocampus, and mesolimbic reward system, underlie addiction pathophysiology. These changes disrupt reward processing, cognitive control, and stress regulation, contributing to heightened cravings and relapse vulnerability. Emerging evidence suggests meditation may modulate these neurobiological pathways.

Objective: This review explores the neurobiological mechanisms underlying addiction and evaluates potential therapeutic effects of meditation on these pathways.

Methods: A narrative review of current literature was conducted, synthesizing evidence from neurobiological studies, clinical trials, metaanalyses, and systematic reviews examining meditation interventions in addiction populations.

Results: Current evidence suggests addiction involves dysregulation of the mesolimbic reward system, with altered dopamine signalling, impaired PFC function, and amygdala hyperactivity. Preliminary research indicates meditation may influence these systems through multiple mechanisms, including modulation of prefrontal-limbic connectivity, stress response normalization via the Hypothalamic-Pituitary-Adrenal (HPA) axis, and enhanced neuroplasticity. However, findings remain mixed, with significant methodological limitations across studies.

Conclusion: While preliminary evidence suggests meditation may engage neurobiological pathways implicated in addiction, current research limitations preclude definitive conclusions. Further rigorous, controlled studies with standardized meditation protocols, larger sample sizes, and longer follow-up periods are essential to establish clinical efficacy and elucidate specific mechanisms of action.

Key Words: *Addiction; Meditation; Neurobiology; Reward system; Prefrontal cortex; Neuroplasticity; Stress regulation; Mindfulness*

¹Department of Yoga, Central University of Rajasthan, Ajmer, Rajasthan-305817, India

²Department of Sangyahan, IMS, Banaras Hindu University, Varanasi, Uttar Pradesh 221005, India

³Department of Sport Psychology, Lakshambai National Institute of Physical Education, NERC, Guwahati, India

⁴Department of Microbiology, Bharathidasan University, Tiruchirappalli, Tamilnadu-620024, India

*Corresponding author: Sanjib Patra, Professor and Head of Department, Department of Yoga, Central University of Rajasthan, Ajmer, Rajasthan-305817, India, E-mail: sanjib.patra@curaj.ac.in

Received: November 30, 2025, Accepted: December 16, 2025, Published: December 23, 2025



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes.

Introduction

Addiction represents a chronic, relapsing condition characterized by compulsive engagement with substances or behaviours, loss of control, and persistence despite adverse consequences. The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) classifies these under “substance-related and addictive disorders,” with diagnostic criteria including intense cravings, tolerance, and withdrawal symptoms [1]. Globally, addiction poses a significant public health burden, with an estimated 39.5 million people suffering from drug use disorders worldwide and addiction-related costs exceeding \$740 billion annually in the United States alone [2,3].

The neurobiological foundations of addiction involve dysfunctions within brain reward circuitry, particularly the mesolimbic dopamine system, Prefrontal Cortex (PFC), and amygdala. These structures regulate reward response, impulse control, emotional regulation, and habit formation. Chronic substance use hijacks this circuitry, producing maladaptive neuroadaptations including heightened sensitivity to addiction-related cues, diminished control over cravings, and increased stress-induced relapse vulnerability [4,5]. Additionally, high rates of comorbid psychological conditions (anxiety, depression, post-traumatic stress disorder) further complicate treatment outcomes [6,7].

Traditional addiction treatments primarily include pharmacological interventions (e.g., methadone, buprenorphine) and behavioral therapies (e.g., cognitive-behavioural therapy, contingency management). While these approaches demonstrate efficacy, relapse rates remain high (40% to 60% within the first year), highlighting the need for complementary therapeutic strategies [8,9].

Within this context, meditation has emerged as a potentially promising adjunctive intervention. Rooted in contemplative traditions, meditation encompasses practices aimed at cultivating mindfulness, self-awareness, and emotional regulation. Preliminary research suggests meditation techniques, including mindfulness meditation, transcendental meditation, and loving-kindness meditation, may modulate neural pathways implicated in addiction [10,11]. Oriented Recovery Enhancement (MORE), such as Mindfulness-Based Relapse Prevention (MBRP) and Mindfulness-Oriented Recovery Enhancement (MORE), have shown preliminary evidence of reducing substance use in some populations, though findings remain inconsistent across studies [12,13].

Neuroimaging studies using Functional Magnetic Resonance Imaging (fMRI) and electroencephalography (EEG) indicate meditation may enhance prefrontal-subcortical connectivity, modulate amygdala reactivity, and influence the Default Mode Network (DMN) [14-16]. However, significant methodological limitations, including small sample sizes, heterogeneous meditation protocols, and lack of active control conditions, limit interpretation of these findings [17].

This review critically examines current evidence regarding meditation’s role in addiction recovery, evaluating both neurobiological mechanisms and clinical outcomes while highlighting methodological limitations and future research needs.

Methodology

Search strategy

A comprehensive literature search was conducted across multiple electronic databases including PubMed/MEDLINE, PsycINFO, Web of Science, and Google Scholar. The search

covered publications from January 2010 through December 2023 to capture recent advances in neuroscience research methodologies. Search terms included combinations of:

- Meditation-related terms: “meditation,” “mindfulness,” “mindfulness-based intervention,” “MBI,” “MBRP,” “MORE,” “transcendental meditation,” “loving-kindness meditation,” “compassion meditation”.
- Addiction-related terms: “addiction,” “substance use disorder,” “SUD,” “substance abuse,” “drug dependence,” “alcohol dependence,” “opioid use disorder,” “relapse,” “craving”.
- Neurobiology-related terms: “neurobiology,” “neuroimaging,” “fMRI,” “functional connectivity,” “dopamine,” “reward system,” “prefrontal cortex,” “neuroplasticity,” “brain changes”.

Studies were prioritized based on:

- methodological rigor (randomized controlled trials, systematic reviews, meta-analyses).
- neurobiological outcomes (neuroimaging findings, neurotransmitter assessments, HPA axis function).
- Clinical relevance to addiction populations.
- Publication in peer-reviewed English-language journals. Emphasis was placed on recent publications (2015-2023) while including seminal foundational studies.

This review synthesizes findings from diverse study designs including randomized controlled trials, neuroimaging investigations, meta-analyses, and systematic reviews. Both positive and negative findings are presented to provide balanced evaluation of current evidence.

Critical limitations of existing research are highlighted throughout, including small sample sizes, heterogeneous meditation protocols, lack of active control conditions, short follow-up periods, and publication bias concerns.

Neurobiology of addiction

Addiction involves complex alterations across multiple brain systems. The mesolimbic reward pathway, comprising the Ventral Tegmental Area (VTA) and Nucleus Accumbens (NAc), plays a central role through dopamine-mediated reward signalling. Chronic substance exposure induces neuroadaptations including dopamine receptor downregulation, reduced dopamine release to natural rewards, and sensitization to drug-associated cues [18].

The prefrontal cortex, particularly dorsolateral and ventromedial regions, mediates executive functions including impulse control, decision-making, and self-regulation. In addition, PFC hypoactivity correlates with impaired behavioral control and increased relapse susceptibility [19]. The amygdala processes emotional salience and stress responses, with hyperactivity contributing to heightened anxiety, negative affect, and cue-induced craving [20]. The hippocampus encodes contextual memories, linking environmental cues with substance use experiences [21].

Key neurotransmitter systems implicated in addiction include dopamine (reward and motivation), glutamate (learning and memory consolidation), GABA (inhibitory control), and endogenous opioids (hedonic tone). The Hypothalamic-Pituitary-Adrenal (HPA) axis dysregulation manifests as elevated basal cortisol and stress hyperreactivity, contributing to negative reinforcement mechanisms underlying continued substance use [22].

Neurobiological Mechanisms of Meditation in Addiction Recovery

Emerging evidence suggests meditation may influence neurobiological pathways implicated in addiction, though mechanisms remain incompletely understood. Preliminary research indicates meditation practice may be associated with structural and functional changes in brain regions central to addiction pathology, including the prefrontal cortex, amygdala, and hippocampus. Neuroimaging studies suggest mindfulness meditation may increase gray matter density in prefrontal regions, potentially enhancing self-regulatory capacity and decision-making processes [23]. Additionally, some research indicates meditation practices may modulate amygdala activity, possibly reducing emotional reactivity to drug-related cues and diminishing the emotional reinforcement of addiction-related stimuli, though findings are not entirely consistent across studies [24,25].

With regard to neurotransmitter systems, preliminary evidence suggests meditation may influence serotonin and dopamine function, neurotransmitters centrally involved in mood regulation and reward processing [26]. By potentially modulating these systems, meditation might help address neurobiological alterations associated with addiction, such as altered dopamine sensitivity and reduced natural reward responsiveness, though the clinical significance and specific mechanisms require further investigation.

While meditation's influence on brain structure, function, and neurochemistry offers promise for addressing addiction-related neurobiological changes, current evidence is limited by small sample sizes, heterogeneous protocols, and insufficient mechanistic understanding. The following sections detail specific proposed mechanisms with critical evaluation of existing evidence (Figures 1 and 2).




Figure 1) Flowchart illustrating the neurobiological pathways influenced by meditation in addiction recovery. Key pathways include regulation of serotonin, dopamine, and neuroplasticity, contributing to improved mood regulation, emotional control, decision-making, and memory, while reducing cravings, relapse risks, and drug cue reinforcement. Brain regions such as the ACC, PFC, PCC, amygdala, and hippocampus are central to these mechanisms. ACC: Anterior Cingulate Cortex; PFC: Pre-frontal Cortex; PCC: Posterior Cingulate Cortex.




Figure 2) Flowchart summarizing neurobiological pathways influenced by meditation in addiction recovery, including modulation of reward pathways, stress regulation, cognitive control, craving-related circuits, and neuroplasticity. LKM: Loving Kindness Meditation; CMT: Compassionate Mind Training; TM: Transcendental Meditation; ANS: Autonomic Nervous System; PFC: Pre-Frontal Cortex.

Modulation of reward pathways

Preliminary neuroimaging studies indicate meditation practice may normalize reward

circuit function. Some research shows increased striatal dopamine release during meditation and enhanced ventral striatum activation to natural rewards following meditation training. However, findings are inconsistent, with other studies reporting no significant dopamine changes. The clinical significance of these alterations for addiction recovery remains unclear [27].

Stress response regulation

Multiple studies suggest meditation may attenuate HPA axis reactivity, reducing stress-induced cortisol release and potentially decreasing relapse vulnerability. Meta-analyses indicate moderate effects of mindfulness interventions on cortisol levels, though heterogeneity across studies is substantial. Long-term practitioners demonstrate lower basal cortisol and attenuated stress responses, though causality cannot be established from cross-sectional comparisons [28].

Enhancement of cognitive control

Neuroimaging evidence suggests meditation training may enhance prefrontal cortex activity and prefrontal-limbic connectivity. Some studies report increased activation in dorsolateral PFC during inhibitory control tasks and strengthened connectivity between PFC and amygdala following mindfulness training. However, effect sizes are generally small to moderate, and clinical translation to improved addiction outcomes remains incompletely demonstrated [29].

Craving-related neural modulation

Preliminary research indicates meditation may reduce activation in craving-related brain regions, particularly the insula and Posterior Cingulate Cortex (PCC). Mindfulness-based interventions show promise in reducing subjective craving intensity and cue-reactivity, though mechanisms remain incompletely

understood. The degree to which neural changes mediate clinical craving reduction requires further investigation [30].

Neuroplasticity and structural changes

Longitudinal studies suggest meditation practice may promote structural brain changes, including increased gray matter density in hippocampus and PFC and alterations in cortical thickness. However, sample sizes in neuroimaging studies are typically small ($n < 50$), limiting generalizability. Whether observed structural changes translate to functional recovery in addiction populations requires systematic investigation [31].

Meditation Practices for Addiction Recovery

Meditation has become a prominent complementary therapy for addiction recovery, due to its ability to address both the neurobiological and psychological factors contributing to substance use disorders. Various meditation techniques have been studied for their effectiveness in reducing addiction severity, promoting emotional regulation, and enhancing cognitive control. Below, we will discuss some of the most commonly utilized meditation practices in addiction treatment, focusing on their specific mechanisms and benefits.

Mindfulness meditation

Mindfulness meditation emphasizes present-moment awareness without judgment [32,33]. Mindfulness-Based Relapse Prevention (MBRP) integrates mindfulness training with cognitive-behavioral relapse prevention strategies [34]. While some trials demonstrate reduced substance use and relapse rates compared to treatment-as-usual, other studies show no significant differences. Meta-analyses suggest small to moderate effects, with substantial heterogeneity across studies

reflecting variations in intervention protocols, control conditions, and outcome measures [35].

Loving-Kindness Meditation (LKM)

Loving-kindness meditation (LKM) cultivates compassion toward self and others [36]. Preliminary evidence suggests LKM may reduce shame, self-criticism, and negative affect factors associated with substance use [37]. However, research specifically examining LKM in addiction populations is limited, with most studies involving non-clinical samples or populations with other mental health conditions [38].

Transcendental Meditation (TM)

Transcendental Meditation (TM) involves silent mantra repetition to achieve restful alertness. Early studies suggested TM might reduce substance use in alcohol-dependent individuals, though methodological limitations (small samples, lack of adequate controls) restrict interpretation [39,40]. More rigorous contemporary research is needed to establish TM efficacy in addiction treatment.

Yoga and meditation

Yoga combines physical postures, breathing exercises, and meditation. Preliminary studies suggest yoga interventions may improve emotional regulation and reduce stress in addiction populations [41,42,43]. However, component analysis distinguishing meditation-specific from exercise-related effects is lacking in most studies.

Discussion

This review synthesized evidence regarding meditation's potential influence on addiction-related neurobiology and clinical outcomes. Preliminary research suggests meditation may modulate multiple neurobiological systems implicated in addiction, including

reward processing, stress regulation, cognitive control, and neuroplasticity. Mindfulness-based interventions show promise for reducing substance use and relapse in some populations, though findings remain inconsistent across studies.

Critical Evaluation and Limitations

Methodological limitations

Current research faces several significant limitations. First, sample sizes in neuroimaging studies are typically small (median $n=30$), limiting statistical power and generalizability. Second, meditation protocols vary substantially across studies in type, duration (ranging from 2 weeks to several months), frequency, and delivery method (group vs. individual, in-person vs. app-based), hindering cross-study comparisons and meta-analytic synthesis. Third, many studies lack active control conditions, raising questions about placebo effects, expectancy, and therapeutic alliance contributions to observed outcomes. Fourth, participant characteristics vary widely, including addiction severity, substance type (alcohol, opioids, stimulants, cannabis), treatment setting (inpatient vs. outpatient), and comorbid conditions, limiting generalizability.

Measurement and Outcome Issues

Outcome measures lack standardization, with studies employing diverse craving scales, substance use measures, and relapse definitions. Neurobiological assessments utilize different imaging protocols, analysis methods, and regions of interest, complicating integration. Follow-up periods are often short (typically 3-6 months), insufficient for assessing sustained recovery given addiction's chronic, relapsing nature. Meditation practice adherence and quality are rarely systematically assessed, obscuring dose-response relationships.

Publication Bias and Reporting

Potential publication bias toward positive findings may inflate apparent effectiveness. Selective outcome reporting within studies and lack of pre-registration for many trials raise concerns about analytical flexibility and p-hacking. Insufficient reporting of adverse events and dropout reasons limits comprehensive risk-benefit assessment.

Theoretical and Mechanistic Considerations

While meditation's effects on attention, emotion regulation, and self-awareness align conceptually with addiction recovery needs, specific mechanisms linking meditation practice to neurobiological changes and clinical outcomes remain incompletely elucidated. Proposed pathways, including enhanced cognitive control, stress reduction, and reward sensitivity normalization, require further validation through mechanistic studies employing mediation analyses, experimental manipulation of specific meditation components, and investigation of individual difference factors predicting treatment response.

Clinical Implications and Future Directions

Despite limitations, existing evidence suggests meditation-based interventions may offer benefits as adjunctive treatments within comprehensive addiction care. However, several research priorities merit attention. Large-scale, multi-site randomized controlled trials with adequate statistical power, active control conditions (e.g., cognitive training, exercise), and extended follow-up periods (≥ 12 months) are essential. Standardization of meditation protocols, including dose, delivery method, and quality assurance procedures, would enhance cross-study comparability.

Component analysis dismantling studies could identify active ingredients within multi-component interventions. Investigation of moderators (addiction severity, substance type, comorbid conditions, demographic factors) and mediators (cognitive, emotional, neurobiological mechanisms) would inform personalized treatment approaches. Integration of multi-modal assessment (neuroimaging, peripheral biomarkers, ecological momentary assessment) would provide comprehensive mechanistic understanding. Finally, pragmatic effectiveness trials in real-world clinical settings would inform implementation and scalability.

Limitations of this Review

This review has several limitations. The narrative synthesis approach, while comprehensive, lacks the quantitative rigor of formal meta-analysis. Exclusion of non-English publications may introduce language bias. The rapidly evolving research landscape means recent studies may not be fully captured. Finally, quality assessment, while considered, was not formally scored using established instruments.

Conclusion

Meditation-based interventions represent a potentially promising complementary approach for addiction recovery, with preliminary evidence suggesting engagement of neurobiological pathways relevant to addiction pathophysiology. However, current research limitations, including small sample sizes, heterogeneous protocols, inadequate control conditions, and insufficient follow-up, preclude definitive conclusions regarding efficacy and mechanisms of action.

While meditation may offer benefits for some individuals within comprehensive addiction treatment, it should not be viewed as a standalone intervention or replacement for evidence-based

treatments. Future research employing rigorous methodologies, standardized protocols, and extended follow-up periods is essential to establish meditation's role in addiction recovery and identify optimal implementation strategies. Until such evidence emerges, meditation-based interventions should be implemented cautiously, with careful monitoring of outcomes and attention to individual patient characteristics and preferences.

Ultimately, addressing the multifaceted challenge of addiction requires integrated approaches combining pharmacological, psychological, social, and potentially contemplative interventions tailored to individual needs. Continued investigation of meditation's neurobiological and clinical effects, alongside development of personalized treatment algorithms, holds promise for enhancing addiction recovery outcomes.

References

1. Klostermann K, Kelley ML. Substance-related and addictive disorders. In: Maddux JE, Winstead BA (eds). *Psychopathology: Foundations for a Contemporary Understanding* (4th edn), Routledge publication, New York, USA. 2016;pp.266-83.
2. United Nation. United Nations office on drugs and crime. <https://www.unodc.org/>.
3. National Center for Drug Abuse Statistics. <https://drugabusestatistics.org/>
4. Koob GF, Volkow ND. Neurobiology of addiction: a neurocircuitry analysis. *Psychiatry*. 2016;3:760-73.
5. Koob GF, Moal ML. Addiction and the brain antireward system. *Annu Rev Psychol*. 2008;59:29-53.
6. Sinha R. Chronic stress, drug use and vulnerability to addiction. *Ann New York Acad Sci*. 2008;1141:105-30.
7. Brady KT, Sinha R. Co-occurring mental and substance use disorders: the neurobiological effects of chronic stress. *Am J Psychiatry*. 2005;162:1483-93.
8. McLellan AT, Lewis DC, O'Brien CP, et al. Drug dependence, a chronic medical illness: implications for treatment, insurance, and outcomes evaluation. *JAMA*. 2000;284:1689-95.
9. Abuse S. Mental Health Services Administration (SAMHSA). 2015 National Survey on Drug Use and Health (NSDUH). Substance Abuse and Mental Health Services Administration. 2020.
10. Tang YY, Tang R, Posner MI. Mindfulness meditation improves emotion regulation and reduces drug abuse. *Drug Alcohol Depend*. 2016;163:S13-8.
11. Garland EL, Froeliger B, Howard MO. Mindfulness training targets neurocognitive mechanisms of addiction at the attention-appraisal-emotion interface. *Front Psychiatry*. 2014;4:173.
12. Bowen S, Chawla N, Grow J, et al. *Mindfulness-Based Relapse Prevention for Addictive Behaviors*. Guilford Publications, New York, USA. 2021.
13. Garland EL, Atchley RM, Hanley AW, et al. Mindfulness-oriented recovery enhancement remediates hedonic dysregulation in opioid users: neural and affective evidence of target engagement. *Sci Adv*. 2019;5:eaax1569.
14. Feldman G, Greeson J, Senville J. Differential effects of mindful breathing, progressive muscle relaxation, and loving-kindness meditation on decentering and negative reactions to repetitive thoughts. *Behav Res Ther*. 2010;48:1002-11.
15. Hölzel BK, Carmody J, Vangel M, et al. Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Res Neuroimaging*. 2011;191:36-43.
16. Brewer JA, Worhunsky PD, Gray JR, et

- al. Meditation experience is associated with differences in default mode network activity and connectivity. *Proc Natl Acad Sci*. 2011;108:20254-9.
17. Brewer JA, Elwafi HM, Davis JH. Craving to quit: psychological models and neurobiological mechanisms of mindfulness training as treatment for addictions. *Psychol Addict Behav*. 2013;27:366-79.
 18. Volkow ND, Fowler JS, Wang GJ. The addicted human brain: insights from imaging studies. *J Clin Investig*. 2003;111:1444-51.
 19. Goldstein RZ, Volkow ND. Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry*. 2002;159:1642-52.
 20. Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacol*. 2010;35:217-38.
 21. LaBar KS, Cabeza R. Cognitive neuroscience of emotional memory. *Nat Rev Neurosci*. 2006;7:54-64.
 22. Volkow N, Li TK. The neuroscience of addiction. *Nat Neurosci*. 2005;8:1429-30.
 23. Lazar SW, Kerr CE, Wasserman RH, et al. Meditation experience is associated with increased cortical thickness. *NeuroReport*. 2005;16:1893-7.
 24. Keng SL, Smoski MJ, Robins CJ. Effects of mindfulness on psychological health: a review of empirical studies. *Clin Psychol Rev*. 2011;31:1041-56.
 25. Davidson RJ, McEwen BS. Social influences on neuroplasticity: stress and interventions to promote well-being. *Nat Neurosci*. 2012;15:689-95.
 26. Zeidan F, Johnson SK, Diamond BJ, et al. Mindfulness meditation improves cognition: evidence of brief mental training. *Conscious Cogn*. 2010;19:597-605.
 27. Brewer JA, Mallik S, Babuscio TA, et al. Mindfulness training for smoking cessation: results from a randomized controlled trial. *Drug Alcohol Depend*. 2011;119:72-80.
 28. Creswell JD, Pacilio LE, Lindsay EK, et al. Brief mindfulness meditation training alters psychological and neuroendocrine responses to social evaluative stress. *Psychoneuroendocrinology*. 2014;44:1-2.
 29. Tang YY, Ma Y, Wang J, et al. Short-term meditation training improves attention and self-regulation. *Proc Natl Acad Sci*. 2007;104:17152-6.
 30. Kober H, Mende-Siedlecki P, Kross EF, et al. Prefrontal-striatal pathway underlies cognitive regulation of craving. *Proc Natl Acad Sci*. 2010;107:14811-6.
 31. Luders E, Toga AW, Lepore N, et al. The underlying anatomical correlates of long-term meditation: larger hippocampal and frontal volumes of gray matter. *NeuroImage*. 2009;45:672-8.
 32. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol Sci Pract*. 2003;10:144-56.
 33. Bowen S, Chawla N, Collins SE, et al. Mindfulness-based relapse prevention for substance use disorders: a pilot efficacy trial. *Subst Abuse*. 2009;30:295-305.
 34. Marlatt GA, Donovan DM. *Relapse Prevention: Maintenance Strategies in the Treatment of Addictive Behaviors*. Guilford Press, New York, USA. 2005.
 35. Witkiewitz K, Marlatt GA, Walker D. Mindfulness-based relapse prevention for alcohol and substance use disorders. *J Cogn Psychother Int Qtrly*. 2005;19:211-28.
 36. Salzberg S, Kabat-Zinn J. *Lovingkindness: The Revolutionary Art of Happiness*. Shambhala Publications, Colorado, United States. 2004.
 37. Keng SL, Smoski MJ, Robins CJ. Effects of mindfulness on psychological health: a review of empirical studies. *Clin Psychol Rev*. 2011;31:1041-56.

38. Galla BM. Within-person changes in mindfulness and self-compassion predict enhanced emotional well-being in healthy, but stressed adolescents. *J Adolesc.* 2016;49:204-17.
39. Haaga DA, Grosswald S, Gaylord-King C, et al. Effects of the transcendental meditation program on substance use among university students. *Cardiol Res Pract.* 2011;2011:537101.
40. Alexander CN, Robinson P, Rainforth M. Treating and preventing alcohol, nicotine, and drug abuse through transcendental meditation: a review and statistical meta-analysis. In: David F, Connell O, Alexander CN (eds). *Self-Recovery*. Routledge Publication, Oxfordshire, UK. 2014;pp.13-87.
41. Van der Kolk B. *The Body Keeps the Score: Brain, Mind, And Body in the Healing of Trauma*. Viking Press, New York, USA. 2014.
42. Chiesa A, Serretti A. Mindfulness-based interventions for chronic pain: a systematic review of the evidence. *J Altern Complement Med.* 2011;17:83-93.
43. Sinha R, Garcia M, Paliwal P, et al. Stress-induced cocaine craving and hypothalamic-pituitary-adrenal responses are predictive of cocaine relapse outcomes. *Arch Gen Psychiatry.* 2006;63:324-31.